

# Orbiter Technical Notes: Nonspherical gravitational field perturbations

Martin Schweiger

September 21, 2005

## 1 Introduction

Orbiter uses a zonal representation of the gravitational potential generated by a celestial body, using a Legendre polynomial series expansion in the latitude  $\theta$ . The perturbations in longitude ( $\phi$ ) are assumed to be negligible. The potential is expressed as

$$U_G(r, \phi, \theta) = -\frac{GM}{r} \left[ 1 - \sum_{n=2}^{\infty} J_n \left( \frac{R}{r} \right)^n P_n(\sin \theta) \right] \quad (1)$$

where  $G$  is the gravitational constant,  $M$  and  $R$  are the mass and mean radius of the central body, respectively,  $r$  is the length of the radius vector,  $J_n$  are the coefficients of the series expansion, and  $P_n$  are the Legendre polynomials of order  $n$ . The first Legendre polynomials are defined as

$$\begin{aligned} P_0(x) &= 1 \\ P_1(x) &= x \\ P_2(x) &= \frac{1}{2}(3x^2 - 1) \\ P_3(x) &= \frac{1}{2}(5x^3 - 3x) \\ P_4(x) &= \frac{1}{8}(35x^4 - 30x^2 + 3) \end{aligned} \quad (2)$$

The acceleration due to the gravitational field of a test mass at point  $\vec{r} = (r, \phi, \theta)$  is then given by the gradient of the potential:

$$\vec{a}_G(r, \phi, \theta) = -\vec{\nabla} U_G(r, \phi, \theta) \quad (3)$$

In spherical polar coordinates, the gradient operator is expressed as

$$\vec{\nabla} = \hat{r} \frac{\partial}{\partial r} + \frac{1}{r} \hat{\theta} \frac{\partial}{\partial \theta} + \frac{1}{r \cos \theta} \hat{\phi} \frac{\partial}{\partial \phi} \quad (4)$$

Substituting equations 1 and 4 into 3 yields

$$\vec{a}_G(r, \phi, \theta) = \hat{r} a_0^{(r)}(r) - \sum_{n=2}^{\infty} \left[ \hat{r} a_n^{(r)}(r, \theta) + \hat{\theta} a_n^{(\theta)}(r, \theta) \right] \quad (5)$$




Figure 1: Planet-relative coordinates and polar unit vectors at a point  $(r, \phi, \theta)$ .

	$J_2$	$J_3$	$J_4$	$J_5$
Mercury	60	-	-	-
Venus	27	-	-	-
Earth	1082.6269	-2.51	-1.60	-0.15
Mars	1964	-	-	-
Jupiter	14750	-	-	-
Saturn	16450	-	-	-
Uranus	12000	-	-	-
Neptune	4000	-	-	-

Table 1: Coefficients ( $\times 10^6$ ) for zonal expansion of planetary gravitational potentials.

with the first terms given by

$$\begin{aligned}
a_0^{(r)}(r) &= -\frac{GM}{r^2} \\
a_2^{(r)}(r, \theta) &= -\frac{3}{2} \frac{GMR^2 J_2}{r^4} (3 \sin^2 \theta - 1) \\
a_2^{(\theta)}(r, \theta) &= 3 \frac{GMR^2 J_2}{r^4} \sin \theta \cos \theta \\
a_3^{(r)}(r, \theta) &= -2 \frac{GMR^3 J_3}{r^5} (5 \sin^3 \theta - 3 \sin \theta) \\
a_3^{(\theta)}(r, \theta) &= \frac{3}{2} \frac{GMR^3 J_3}{r^5} (5 \sin^2 \theta \cos \theta - \cos \theta) \\
a_4^{(r)}(r, \theta) &= -\frac{5}{8} \frac{GMR^4 J_4}{r^6} (35 \sin^4 \theta - 30 \sin^2 \theta + 3) \\
a_4^{(\theta)}(r, \theta) &= \frac{5}{2} \frac{GMR^4 J_4}{r^6} (7 \sin^3 \theta \cos \theta - 3 \sin \theta \cos \theta)
\end{aligned} \tag{6}$$

The coefficients  $J_n$  used by Orbiter are listed in Table 1.

The field perturbations can lead to a rotation of the orbit trajectory of a satellite. This rotation can be expressed in terms of the movement of the longitude of the ascending node ( $\Omega$ ) and the movement of the argument of periapsis ( $\omega$ ). If only terms up

to  $J_2$  are included, approximate values of the movements  $\partial\Omega/\partial t$  and  $\partial\omega/\partial t$  are given by

$$\frac{\partial\Omega}{\partial t} = -\frac{3n}{2} \left(\frac{R}{a}\right)^2 \frac{\cos i}{(1-e^2)^2} J_2 \quad (7)$$

$$\frac{\partial\omega}{\partial t} = \frac{3n}{4} \left(\frac{R}{a}\right)^2 \frac{5\cos^2 i - 1}{(1-e^2)^2} J_2 \quad (8)$$

where  $n = 2\pi/P$  is the mean motion (with orbit period  $P$ ),  $a$  is the mean distance,  $e$  is the eccentricity, and  $i$  is the inclination.

### Example: calculate the inclination for a sun-synchronous polar orbit

A sun-synchronous orbit exploits the propagation of the line of nodes to keep the orbital plane synchronised with the relative position of the sun. A satellite can for example be placed in a sun-synchronous orbit so that it continuously flies over the planet's terminator line. From 7 we have

$$\cos i = -\frac{2}{3n} \left(\frac{a}{R}\right)^2 \frac{(1-e^2)^2}{J_2} \frac{\partial\Omega}{\partial t} \quad (9)$$

A sun-synchronous orbit requires the line of nodes to move at a rate of  $2\pi$  per year. For Earth, this is equivalent to  $\partial\Omega/\partial t = 1.99 \cdot 10^{-7}$  rad/s (about 0.99 deg. per day). Assume a circular orbit ( $e = 0$ ) at an altitude of 300 km ( $a = 6671010$  m, with  $R_E = 6371010$  m). With  $P = 2\pi\sqrt{a^3/\mu_E}$  we get  $n = \sqrt{\mu_E/a^3} = 0.0012$  rad/s. This leads to

$$\cos i_{\text{sync}} = -\frac{2}{0.0035} \left(\frac{6678137}{6378137}\right)^2 \frac{1.99 \cdot 10^{-7}}{0.001082630} = -0.116, \quad (10)$$

or  $i_{\text{sync}} = 96.7$  deg.